

Avian influenza (bird flu)

In the guide

[Avian influenza prevention measures](#)

[Possible impact of the disease](#)

[Clinical signs](#)

[Procedures when disease is confirmed](#)

[Biosecurity](#)

[Legislation applicable to avian influenza](#)

[Effect on food and humans](#)

[Moving eggs](#)

[Disease alerts](#)

[Trading Standards](#)

[In this update](#)

[Key legislation](#)

This guidance is for Wales

Avian influenza is a highly infectious viral disease affecting the respiratory, digestive and/or nervous system of birds.

The severity depends upon the strain of the virus and the type of bird infected. Some strains (known as 'highly pathogenic avian influenza') can cause severe disease in poultry, with a high death rate. The disease can develop so rapidly that birds may die without showing any previous signs of disease.

Other strains (known as 'low pathogenic avian influenza') usually result in milder, less significant disease. However, certain low pathogen strains can mutate into highly pathogenic strains.

Avian influenza prevention measures

Information on the latest situation across Great Britain can be found on [GOV.UK](#), [GOV.SCOT](#) and [GOV.WALES](#). See also the Animal and Plant Health Agency's [interactive map](#).

Possible impact of the disease

Highly pathogenic avian influenza can cause severe disease in susceptible birds. Low pathogenic avian influenza generally causes mild disease or no disease at all.

All bird species are thought to be susceptible to avian influenza. Migratory birds such as wild ducks and geese can carry the viruses, often without symptoms of illness, and show the greatest resistance to

infection. However, domestic poultry flocks are particularly vulnerable to epidemics of a rapid, severe and fatal form of the disease.

Legislation and processes for management of the disease are different, depending on the circumstances of the outbreak - for example, whether the high pathogenic strain H5N1 is confirmed in poultry, other captive birds or wild birds, and whether the disease occurs at a farm, slaughterhouse or border inspection post.

The highly pathogenic H5N1 strain has shown the ability to jump the species barrier occasionally and cause severe disease in humans. It has not shown the ability to move easily between humans, but this is a cause for concern (see below).

Clinical signs

The severity depends upon the strain of virus and the type of bird infected. Birds infected with highly pathogenic avian influenza may die suddenly or show a range of clinical signs.

Individual birds with highly pathogenic avian influenza may show the following symptoms:

- Nervous signs (tremors and incoordination)
- Depression, sneezing and coughing
- Swollen, congested and oedematous wattles
- Haemorrhages on the hock (lower part of leg)
- Diarrhoea

In the flock, symptoms can be:

- Unusual quietness, decreased activity levels
- Decreased levels of vocalisation
- Decreased levels of feed and water consumption
- Decreased egg production

It is important to note that when infection is due to a highly pathogenic form of avian influenza, the onset of clinical signs is sudden, severe, short lasting and mortality is extremely high (sometimes 100%).

Avian influenza is a notifiable disease. If you suspect any type of bird flu, you must tell the Animal and Plant Health Agency (APHA) immediately by telephoning 0300 303 8268. Failure to do so is an offence.

Procedures when disease is confirmed

The premises where disease is confirmed will be put under restriction so poultry or other captive birds are isolated and cannot move on or off. It will be referred to as the infected premises. An approved disinfectant must be used to disinfect footwear, clothing and vehicles before entering or leaving the premises.

A protection zone of 3 km and a surveillance zone of 10 km may be put in place around the infected premises where the disease has been confirmed. There are certain restrictions for keepers of poultry that are within the protection and surveillance zones. Keepers can now apply for a licence to move poultry from a restricted zone, using APHA's [online animal disease licensing service](#).

If avian influenza is confirmed, it will be controlled in line with the Welsh Government [contingency plan for exotic animal diseases](#) and the [notifiable avian disease control strategy](#) will be implemented.

Biosecurity

Biosecurity measures should be practised routinely. Avian influenza is spread through bird-to-bird contact and indirectly through contaminated feed, water, equipment, etc.

The boots, clothing and hands of any person who has been in contact with infected animals can spread the disease.

The Welsh Government has published a poster [on protecting your birds from avian influenza](#). There is also guidance on [how to spot and report the disease](#).

Although the risk to kept birds has reduced, all bird keepers are advised to complete the [biosecurity self-assessment checklist](#). This should be completed annually and kept as part of the flock records. This is a legal requirement when an avian influenza prevention zone (AIPZ) has been declared in Wales.

Legislation applicable to avian influenza

The Animal Health Act 1981 provides powers for the control of outbreaks of avian influenza. It was amended in 2002 to provide more powers to deal with foot-and-mouth disease and these powers were extended by the Avian Influenza and Newcastle Disease (England and Wales) Order 2003, so that they are now exercisable in relation to avian influenza (and Newcastle disease).

The Act provides for the slaughter of diseased poultry, poultry suspected of disease, poultry exposed to disease and poultry that the Welsh Government thinks should be slaughtered to prevent the spread of disease (a 'firebreak' cull).

The Diseases of Poultry (Wales) Order 2003 introduced a number of key amendments to the Animal Health Act 1981. The definition of poultry within section 87(4) of the Animal Health Act 1981 was amended to include all birds (including those in captivity) and the definition of disease in section 88(3) was extended to include all diseases of birds.

The Avian Influenza and Influenza of Avian Origin in Mammals (Wales) (No 2) Order 2006 allows for a flexible and risk-based approach to disease control, which will enable industry to continue to operate in a safe and biosecure manner.

The Order contains preventive measures such as:

- Power for the Welsh Government to declare an avian influenza prevention zone
- Ban or limitation of bird gatherings
- Separation of birds from wild birds
- Surveillance for avian influenza

The Order includes a range of provisions for application in cases of suspected and confirmed avian influenza, both highly pathogenic and low pathogenic.

The Avian Influenza (H5N1 in Wild Birds) (Wales) Order 2006 provides controls in the event that avian influenza H5N1 is found in wild birds.

The Avian Influenza (Vaccination) (Wales) Regulations 2006 prohibit vaccination but allow the Welsh Government to declare an emergency vaccination zone or a preventive vaccination zone. The Welsh Government can also serve an emergency vaccination notice on individual premises. Within the zones or on individual premises, the Welsh Government can require the vaccination of poultry and other captive

birds. Movement restrictions will apply within the zones.

The Avian Influenza (Preventive Measures) (Wales) Regulations 2006 ban fairs, markets, shows or other gatherings of poultry or birds. In practical terms, bird gatherings have ordinarily been permitted to take place under a general licence, but the organiser must notify APHA on 0300 303 8268 or customeradvice@apha.gov.uk.

The current position in relation to bird gatherings is that gatherings of all poultry are no longer permitted in Wales. Gatherings of captive birds may still take place as long as you meet the requirements of the [captive birds gathering general licence](#).

Keepers of poultry and other captive birds are required to register their flocks with APHA. See '[Registration and records of poultry and other captive birds](#)' for more information on registration requirements.

The Regulations also restrict the vaccination of birds in zoos. The Welsh Government can require vaccination of birds in zoos to be carried out. Certain measures apply to vaccinated birds in zoos, and the Welsh Government can require surveillance at these zoos.

The Avian Influenza and Newcastle Disease (Contingency Planning) (Wales) Order 2005 provides for the preparation of a national contingency plan indicating the arrangements to be put in place for the control and eradication of avian influenza (and Newcastle disease). The national contingency plans for these diseases must be issued on an annual basis for public consultation and issued to Parliament.

Effect on food and humans

[Advice from the World Health Organisation](#) is that there is no health risk from well cooked poultry meat or from eggs.

Avian influenza is primarily a disease of birds. Humans can only be infected with the disease through close contact with live infected birds.

Transmission of avian influenza viruses to people remains relatively rare, and in most cases occurs as a result of direct contact with infected poultry or other birds, or their faeces. At present, the virus cannot transfer directly from human to human.

There is concern that the virus may change (or mutate) to emerge as a new virus that is transmissible between people and capable of causing disease in people, birds and other animals.

Public health control measures in any outbreak of avian influenza amongst poultry will therefore aim to protect people against avian influenza and also protect against the risk of any mutation of the virus.

For more information please see the section of the GOV.UK website on [avian influenza: guidance, data and analysis](#).

Moving eggs

If you want to move table eggs into, within or out of a protection or surveillance zone to an egg processing plant, you will need to comply with the relevant law, conditions and instructions associated with the [general licence EXD541\(HPAI\)\(GB\)](#), information on which can be found on the GOV.UK website.

This general licence is used for movements of table eggs during outbreaks of 'H5N' strains of avian

influenza - for example, H5N1 and H5N5. You must not move table eggs into, within or out of a protection or surveillance zone during outbreaks of any other strains of HPAI.

Disease alerts

Livestock keepers can stay up to date with the latest avian influenza developments via the APHA [alert subscription service](#).

Trading Standards

For more information on the work of Trading Standards services and the possible consequences of not abiding by the law, please see '[Trading Standards: powers, enforcement and penalties](#)'.

In this update

New section: 'Moving eggs'.

Last reviewed / updated: May 2026

Key legislation

- [Animal Health Act 1981](#)
- [Diseases of Poultry \(Wales\) Order 2003](#)
- [Avian Influenza and Newcastle Disease \(England and Wales\) Order 2003](#)
- [Avian Influenza and Newcastle Disease \(Contingency Planning\) \(Wales\) Order 2005](#)
- [Avian Influenza \(Preventive Measures\) \(Wales\) Regulations 2006](#)
- [Avian Influenza and Influenza of Avian Origin in Mammals \(Wales\) Order 2006](#)
- [Avian Influenza and Influenza of Avian Origin in Mammals \(Wales\) \(No 2\) Order 2006](#)
- [Avian Influenza \(Vaccination\) \(Wales\) Regulations 2006](#)
- [Avian Influenza \(H5N1 in Wild Birds\) \(Wales\) Order 2006](#)

Please note

This information is intended for guidance; only the courts can give an authoritative interpretation of the law.

The guide's 'Key legislation' links go to the legislation.gov.uk website. The site usually updates the legislation to include any amendments made to it. However, this is not always the case. Information on all changes made to legislation can be found by following the above links and clicking on the 'More Resources' tab.

© 2026 Chartered Trading Standards Institute

Source URL:

<https://www.businesscompanion.info/en/quick-guides/animals-and-agriculture/avian-influenza-bird-flu-w>